



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2014

Applying 'Patient Blood Management' in the trauma center

Theusinger, Oliver M ; Stein, Philipp ; Spahn, Donat R

Abstract: PURPOSE OF REVIEW: The purpose of this review is to highlight the use of tranexamic acid, point-of-care testing, algorithm-based treatment of trauma-associated coagulopathy with factor concentrates to reduce blood loss and transfusion requirements in order to improve outcome. In addition, the management of patients on new oral anticoagulants, drugs with renewed interest and the tolerance of relatively low hemoglobin levels in the context of trauma will be discussed. **RECENT FINDINGS:** Early administration of tranexamic acid reduces mortality without increasing the risk of thromboembolic events. Point-of-care testing is increasingly recommended. Goal-directed individualized coagulation algorithms with the use of factor concentrates allow reducing the amount of allogeneic blood products to be administered. Treatment of trauma patients with one of the new oral anticoagulants is challenging. Furthermore, new mechanisms have been discovered such as deep neuromuscular blockade to better tolerate acute anemia. **SUMMARY:** Applying Patient Blood Management concept to the trauma patient is possible and efficacious. Antihyperfibrinolytics such as tranexamic acid, point-of-care testing and coagulation algorithms with the use of factor concentrates allow a reduction of the number of transfusions, the costs and will likely ameliorate outcome of major trauma patients.

DOI: <https://doi.org/10.1097/ACO.0000000000000041>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-95793>

Journal Article

Published Version

Originally published at:

Theusinger, Oliver M; Stein, Philipp; Spahn, Donat R (2014). Applying 'Patient Blood Management' in the trauma center. *Current Opinion in Anaesthesiology*, 27(2):225-232.

DOI: <https://doi.org/10.1097/ACO.0000000000000041>



Applying 'Patient Blood Management' in the trauma center

Oliver M. Theusinger, Philipp Stein, and Donat R. Spahn

Purpose of review

The purpose of this review is to highlight the use of tranexamic acid, point-of-care testing, algorithm-based treatment of trauma-associated coagulopathy with factor concentrates to reduce blood loss and transfusion requirements in order to improve outcome. In addition, the management of patients on new oral anticoagulants, drugs with renewed interest and the tolerance of relatively low hemoglobin levels in the context of trauma will be discussed.

Recent findings

Early administration of tranexamic acid reduces mortality without increasing the risk of thromboembolic events. Point-of-care testing is increasingly recommended. Goal-directed individualized coagulation algorithms with the use of factor concentrates allow reducing the amount of allogeneic blood products to be administered. Treatment of trauma patients with one of the new oral anticoagulants is challenging. Furthermore, new mechanisms have been discovered such as deep neuromuscular blockade to better tolerate acute anemia.

Summary

Applying Patient Blood Management concept to the trauma patient is possible and efficacious. Antifibrinolytics such as tranexamic acid, point-of-care testing and coagulation algorithms with the use of factor concentrates allow a reduction of the number of transfusions, the costs and will likely ameliorate outcome of major trauma patients.

Keywords

goal-directed transfusions, Patient Blood Management, ROTEM, thromboelastometry, transfusion management, traumatic coagulopathy

INTRODUCTION

Trauma-associated bleeding leading to death has been reported in up to 40% of all casualties in civilian trauma centers [1,2,3^{***}]. Furthermore, it is assumed that up to one-quarter of trauma-associated deaths might be preventable if early and rapid control of blood loss and coagulopathy is established [4,5^{***}]. Injuries leading to exsanguination are to be located in the majority of the cases in the thorax, abdomen and pelvis [6]. In addition, prognosis of these patients is worse when coagulopathy is present at hospital admission [7–11]. For this reason, early, aggressive, and rapid hemostatic interventions are the key to prevent exsanguination and avoid or minimize massive transfusion requirements [3^{***}].

Since 2010 the World Health Organization urges member states to utilize transfusion alternatives and develop individualized Patient Blood Management Programs in order to reduce transfusion needs.

The three pillars of Patient Blood Management consist of: 'detection and treatment of preoperative anemia, reduction in perioperative red blood cell loss, and harnessing and optimizing the patient-specific physiological reserve of anemia (including restrictive hemoglobin transfusion triggers)' [12^{*,}13^{*,}14^{***},15^{*,}16^{***}].

The first pillar is obviously not applicable in the context of trauma, but the second and third pillars are to be used also in the case of trauma. Standard coagulation tests are used widely to

Institute of Anesthesiology, University and University Hospital of Zurich, Zurich, Switzerland

Correspondence to Oliver M. Theusinger, MD, Institute of Anesthesiology, University Hospital and University of Zurich, Rämistrasse 100, Zurich CH-8091, Switzerland. Tel: +41 44 255 27 10; fax: +41 44 255 44 09; e-mail: oliver.theusinger@usz.ch

Curr Opin Anesthesiol 2014, 27:225–232

DOI:10.1097/ACO.000000000000041

KEY POINTS

- The second and third pillar of Patient Blood Management Programs can be applied in the setting of trauma patients to reduce the use of blood products.
- The use of new oral anticoagulants is a new challenge in trauma, as reversal of these drugs is difficult and partially impossible.
- Tranexamic acid is becoming one of the key drugs to reduce bleeding and reduce mortality in trauma patients.

identify trauma-induced coagulopathy but the value of these tests has been challenged in recent years. Routine coagulation tests were not developed to identify bleeding in trauma. Furthermore these tests are neither predictive for bleeding nor validated for use in patients with major trauma [17]. For this reason point-of-care devices, such as rotational thromboelastometry (ROTEM, TEM Innovations GmbH, Munich, Germany) or thromboelastography (TEG, Haemonetics Corp, Niles, IL, USA), are used increasingly to treat bleeding in trauma and are actually recommended to assist in characterizing the coagulopathy and guide the hemostatic therapy [3^{••},18–21,22^{••}].

The purpose of this review is to highlight the use of tranexamic acid, point-of-care testing, algorithm-based treatment of trauma-associated coagulopathy with factor concentrates to reduce blood loss and transfusion requirements in order to improve outcome. In addition, the management of patients on new oral anticoagulants, drugs with renewed interest and the tolerance of relatively low hemoglobin levels in the context of trauma will be discussed.

TRANEXAMIC ACID

In 1950, Okamoto *et al.* [23] described for the first time the lysin derivate tranexamic acid (trans-4-aminomethylcyclohexane-1-carboxylic acid, TXA) that inhibits the action of plasmin. The use of TXA showed a significant blood sparing effect in elective surgery [24]. In 2010, the CRASH-2 Study was the first randomized, placebo-controlled trial to investigate TXA use in trauma. Application of TXA to adult trauma patients in the emergency department reduced all-cause mortality from 16.0 to 14.5% ($P=0.0035$) and the risk of death due to blood loss from 5.7 to 4.9% ($P=0.0077$). The dosing regimen was 1 g intravenously (i.v.) over 10 min followed by an infusion of 1 g over 8 h. Expected severe side-effects, in particular thrombotic events

such as myocardial infarction and vascular occlusive events were specifically assessed and found to occur less frequently in the tranexamic acid group [25]. This study led to a worldwide increased scientific interest for this antifibrinolytic drug (2009: 84 PubMed listed publications, January–October 2013: 229 publications. Keyword: tranexamic acid).

Recent studies review dose, timing of application, exact indication and potential contraindications or adverse events of TXA use in trauma patients. Grassin-Delyle *et al.* [26[•]] proposed a dosing scheme of TXA based upon a pharmacokinetic open two-compartment model with linear elimination. This implicates bolus application with continuous infusion of TXA. The validity for trauma patients may be limited as the calculations were made with children on cardiopulmonary bypass. A systematic review of 104 randomized trials found that the highly significant reduction of surgical blood loss was largely independent of the i.v. TXA doses from 5.5 to 300 mg/kg. The authors conclude that a total dose of 1 g is sufficient for most adult patients [27,28^{••}].

In the prehospital setting 40 trauma patients received TXA: 55% of the victims had penetrating injuries, 45% had blunt trauma. The authors concluded that prehospital administration of TXA at the site of injury is feasible and that no direct adverse drug events were observed [29^{••}]. However, the study design and sample size did not allow assessing patient's outcome [29^{••}]. The re-analysis of the CRASH-2 data showed that early TXA administration within 1 h of arrival in the emergency room significantly reduced the risk of death due to bleeding from 7.7 to 5.3% and from 6.1 to 4.8% when given within 1–3 h. Later administration resulted in an increased mortality due to bleeding (4.4 vs. 3.1%) [30]. Moreover, the use of TXA has been shown to be highly cost-effective [31^{••}].

TXA had also been successfully introduced in military treatment algorithms. Despite greater Injury Severity Scores after suffering combat injuries, mortality was lowest for patients receiving cryoprecipitate and TXA (11.6%, $n=168$) and TXA only (18.2%, $n=148$) compared with cryoprecipitate alone (21.4%, $n=258$) or no cryoprecipitate/TXA (23.6%, $n=758$). In addition, a systematic Cochrane review provides evidence that TXA reduces blood transfusions in patients requiring emergency or urgent surgery [32[•]].

On the basis of WHO data and systematic literature review, Ker *et al.* [28^{••}] estimated that approximately 400 000 trauma patients die in hospital due to bleeding. They calculated that approximately 128 000 deaths might be averted if these patients received TXA within 1–3 h of injury.

In accordance with the European trauma treatment guideline for the management of bleeding and coagulopathy following major trauma, 'we emphasize that the first dose of TXA should be given to all patients with trauma and significant bleeding prehospital at the scene of accident or at the latest within 3 h after the initial trauma' [3^{''}].

POINT-OF-CARE TESTING

In trauma, coagulopathy is common and multifactorial. Early recognition and treatment is likely to reduce blood loss, the use of blood products, and morbidity and mortality [3^{''},33,34]. In recent publications, transfusion algorithms guided goal-directed transfusion (Fig. 1) therapy based on laboratory variables and point-of-care testing were described [3^{''},35]. Thromboelastography and thromboelastometry both measure and graphically display the changes in viscoelasticity at all stages of the developing and resolving clot, and provide the first results within 5–10 min, whereas laboratory values take from 30 min and up to 90 min delaying treatment in patients [36]. The method of the ROTEM has been described in detail elsewhere [37].

To monitor coagulation in the trauma patient, various tests are available including International Normalized Ratio, aPTT, thrombin time, fibrinogen testing (often by the Clauss method) [38], platelet count, platelet function testing, and factor XIII and FVIII determination. Proof-of-concept monitoring of blood coagulation at the bedside is not only desirable, but is becoming increasingly recommended [3^{''}].

ACUTE ANEMIA

In contrast to routine cases, trauma patients undergoing surgery are generally not accessible for pre-operative correction of anemia. However, severe anemia (hemoglobin as low as 14 g/l) can be survived without squeal, even without allogeneic blood transfusion [39]. Nevertheless, the European trauma treatment guidelines recommend a hemoglobin target range of 70–90 g/dl [3^{''}].

In a recent trial, pigs were randomized into three different groups and hemodiluted with hydroxyethyl starch comparing no hemodilution vs. hemoglobin of 4.0 g/dl and vs. the critical hemoglobin level of 2.7 g/dl. In the hemodiluted state, 10 mg/kg of pimonidazole was injected, which forms protein adducts in hypoxic cells. Interestingly, metabolic parameters and oxygen consumption did not show that tissue oxygenation was restricted before reaching a hemoglobin level of 2.7 g/dl. Kidneys and

skeletal muscle showed enhanced pimonidazole binding and vascular endothelial growth factor expression at a hemoglobin level of 4 g/dl. Other organs like heart, brain and liver showed no signs of tissue hypoxia at hemoglobin levels of 4 g/dl. Acute anemia tolerance is thus organ specific and needs to be further elucidated [40^{''}].

Another interesting finding was made by Pape *et al.* [41^{''}] showing that deep neuromuscular blockade increases the tolerance of acute normovolemic anemia. The possible mechanism seems to involve a reduction of skeletal muscular oxygen consumption for the benefit of vital organs. Neuromuscular blockade therefore might play an important role in the treatment of trauma victims as critical levels of hemoglobin may be better tolerated.

COAGULATION FACTOR CONCENTRATES

The transfusion of fresh frozen plasma (FFP) leads to adverse effects such as an increased mortality, multiple organ failure, risk of infection, lung injury and immunomodulation [42,43^{''}]. The effectiveness of FFP compared with fibrinogen concentrate regarding clinical endpoints as blood loss, transfusion requirements, hospital length of stay, survival and plasma fibrinogen concentration is favorable for fibrinogen concentrates [44]. In patients with massive transfusion, the use of FFP or fibrinogen is indicated according to the European trauma Guidelines. If FFP is used, a plasma : red blood cell ratio of least 1 : 2 is suggested. However, plasma transfusions are to be avoided in patients without substantial bleeding [3^{''}].

In many bleeding situations, including trauma, fibrinogen is the first coagulation factor to become critically low. Early testing for fibrinogen concentration and specially for fibrinogen activity has been proposed and implemented in many centers [3^{''},35,45–48].

In addition, target levels of fibrinogen concentrations for trauma patients have been defined at 1.5–2.0 g/l in the European Trauma Treatment Guidelines [3^{''}]. As fibrinogen is a central element of hemostasis, achieving these target levels is important. Administering fibrinogen concentrates is advantageous over FFP transfusion because the fibrinogen concentration in FFP is highly variable and rather low with 2 g/l on average [49]. Therefore, large volume transfusions of FFP would be necessary to effectively increase the fibrinogen concentration, and achieving the target fibrinogen concentration of 1.5–2.0 g/l is virtually impossible given the average fibrinogen concentration of 2 g/l. When replacing most FFP transfusions with fibrinogen


Diagnostic	Intervention
Preoperative history <ol style="list-style-type: none"> Drugs affecting coagulation <ul style="list-style-type: none"> Antiplatelet drugs Heparin Oral anticoagulation (Vit. K antagonists), Xa antagonists, IIa antagonists Coagulation status? HIT II? 	ROTEM after anesthesia induction <ul style="list-style-type: none"> Transplant surgery Cardiac and vascular surgery Difficult cancer surgery Liver insufficiency Intra-abdominal sepsis Emergency room entry
Blood loss > 50% with diffuse bleeding	
ROTEM analysis <ul style="list-style-type: none"> EXTEM, INTEM, FIBTEM, APTM HEPTEM in heart and vascular surgery 	Target values <ul style="list-style-type: none"> Normothermia (temp > 35°C) Normocalcaemia (Ca > 1.15 mmol/l) No acidosis (pH > 7.2) Hematocrit > 0.21 Hypotension (MAP 55–60 mmHg) Crystalloid and/or colloid volume substitution
FIBTEM < 7 mm	Fibrinogen 2–4 g i.v. (maximal 3x2g), after a total of 6 g give FXIII
INTEM (CT and CFT prolonged) and HEPTEM normal or ACT pathological and heparinase ACT normal	Protamine sulfate 1:1 to heparin crystalloid and colloid volume substitution
EXTEM/INTEM : Decrease of MCF after maximum was reached APTEM : normal 	Tranexamic acid <ul style="list-style-type: none"> 15 mg/kg body weight as bolus iv 1–2 mg/kg/h during surgery iv as perfusion
Ongoing diffuse bleeding	
EXTEM/INTEM MCF < 40 mm CT EXTEM/INTEM normal MCF FIBTEM < 7 mm Hematocrit > 0.21 MCF FIBTEM > 7 mm Platelets < 50 000/ l (< 100 000/ l in cardiac surgery or in patients suffering from traumatic brain injury) Coagulation test incl. F XIII, F V, INR, PT, aPTT	Fibrinogen up to 6 g, followed by Factor XIII 15 U/kg body weight crystalloid and colloid volume substitution Platelet concentrates Target of Factor XIII: > 60% (Factor XIII 15 U/kg body weight) Target of Factor V: > 20% (in particular in liver insufficiency /trauma or intra-abdominal sepsis: 2–4 U FFP)
Ongoing diffuse bleeding	
Quick's value < 30% and Factor V > 20% or EXTEM/INTEM: CT, CFT prolonged	4 factor prothrombin complex 1000–2000 IU <ul style="list-style-type: none"> Factor II, VII, IX and X Depending on the patients' body weight
In case of massive transfusion	Target hematocrit 0.21–0.24
If massive diffuse bleeding continues and	
Treated acidosis Treated hypothermia Treated hypocalcaemia Hematocrit: 0.21–0.24 Excluded DIC Fibrinogen was substituted Platelets > 50 000/ l (> 100 000/ l in cardiac surgery or in patients suffering from traumatic brain injury)	Recombinant factor VIIa 60 g/kg body weight A second dose of 60 g/kg body weight can be given again after 2–4 hours, if bleeding has not completely stopped.

FIGURE 1. Third version of the transfusion algorithm of the University Hospital of Zürich 2013, Switzerland. BW = body weight (modified according to [35], with permission of Lippincott Williams & Wilkins).

concentrate, monitoring factor XIII levels is advisable after the replacement of approximately 50% of the blood volume; in addition, a 60% factor XIII activity level is to be maintained with the administration of factor XIII concentrate [35,50–52].

Apart from single-factor concentrates, prothrombin complex concentrates (PCC) are also part of factor-based algorithms [35,46–48,53,54]. Individual PCCs differ regarding their factors contained, their relative composition, and their thrombotic potential. In general, PCCs are used for the reversal of oral

anticoagulants [55]. The use of PCC is also suggested by the update of the European Trauma Guidelines in 2013 if the initiation of clot formation is prolonged in ROTEM [3[■]]. PCC should be used only within strict algorithms, the dose should be small and repeat doses should be given with caution to minimize thrombotic risks [54].

Actual studies indicate that a high amount of RBCs, FFPs and platelets can be saved without additional risks for patients by using a coagulation factor concentrate-based coagulopathy management in trauma [56]. Based upon data from four European countries (UK, Germany, Italy and Switzerland), blood substitution and blood products were calculated and found to account for approximately 27% of all costs associated with trauma care on the ICU [57]. The reduced frequency of septic complications and organ failure observed here that translated into trends toward reduced days on ventilator while on ICU and shorter overall in-hospital length of stays may also contribute to cost reduction in acute trauma care without increasing the risk for the individual patient.

DRUGS WITH RENEWED INTEREST

Desmopressin (DDAVP: 1-deamino-8-D-arginine vasopressin, 0.3 mcg/kg) enhances platelet adherence and is the first choice in the treatment of bleeding patients with quantitative deficiency (type 1) and some with qualitative defects (type 2) of von Willebrand factor [58,59]. Patients treated with aspirin, ADP receptor inhibitors such as clopidogrel could benefit from point-of-care testing to assess the individual efficacy of desmopressin treatment [60]. Furthermore, the European Trauma Guidelines suggest that patients treated with aspirin only may benefit from a single dose of desmopressin [3[■]]. Should the treatment with desmopressin be insufficient, transfusing platelets is the ultimate option. Other antiplatelet drugs such as ticagrelor might not be easily reversed with desmopressin and even platelet transfusion may be relatively ineffective [3[■],61[■]].

NEW DRUGS

New oral anticoagulants acting as direct thrombin (dabigatran) and factor Xa inhibitors (apixaban and rivaroxaban) are now encountered also in the bleeding trauma patient. Experience and treatment options are limited.

There are reports of continuous bleeding after trauma, in emergency surgery and in spontaneous intracerebral hemorrhage [62[■]–64[■]]. This is probably

not because of the higher risk of bleeding *per se* but to the persistence of effective plasma levels of the anti-coagulatory substances [65[■]].

In healthy volunteers, a high dose (50 U/kg) of nonactivated PCC was able to completely reverse antifactor Xa effects on laboratory parameters due to rivaroxaban [66].

The use of activated factor VII to treat bleeding under new oral anticoagulants is off label. Only ex-vivo and in-vitro studies exist, showing theoretical and limited effect [67,68[■]] with case reports of successful hemostatic treatment [69]. Hemodialysis was reported to be effective for dabigatran removal [70[■],71]. Because of the high volume of distribution, Chang *et al.* [72], however, saw a rebound of dabigatran plasma levels after discontinuation of dialysis.

Convincing clinical evidence for the reversal of action of the new oral anticoagulants is lacking; no studies evaluating the clinical success and the possible thrombotic adverse events exist to date. Immediate discontinuation of the anticoagulant is mandatory. Activated charcoal is indicated for the absorption of dabigatran within 2 h after ingestion [73].

Specific antagonists for factor Xa mediated anticoagulation are needed for immediate reversal in the bleeding patient. R-Antidote (PRT064445) may be a potential candidate and is currently under investigation [74].

CONCLUSION

Applying the second pillar of Patient Blood Management to the trauma patient is possible and highly efficacious. Antihyperfibrinolytics such as tranexamic acid, point-of-care testing and coagulation algorithms with the use of factor concentrates allow a reduction of the number of transfusions, the costs and will likely ameliorate outcome of major trauma patients.

Acknowledgements

None.

Conflicts of interest

O.M.T. has received honoraria or travel support for consulting or lecturing from the following companies: CSL Behring Schweiz, Zurich, Switzerland, Vifor SA, Villars-sur-Glâne, Switzerland, Roche Pharma (Schweiz) AG, Reinach, Switzerland, Pentapharm AG, München, Germany, TEM International, München, Germany.

D.R.S. academic department is receiving grant support from the Swiss National Science Foundation, Berne, Switzerland (grant numbers: 33CM30_124117 and 406440-131268), the Swiss Society of Anesthesiology

and Reanimation (SGAR), Berne, Switzerland (no grant numbers are attributed), the Swiss Foundation for Anesthesia Research, Zurich, Switzerland (no grant numbers are attributed), Bundesprogramm Chancengleichheit, Berne, Switzerland (no grant numbers are attributed), CSL Behring, Berne, Switzerland (no grant numbers are attributed), Vifor SA, Villars-sur-Glâne, Switzerland (no grant numbers are attributed).

D.R.S. was the chairman of the ABC Faculty and a member of the ABC Trauma Faculty which both are managed by Thomson Physicians World GmbH, Mannheim, Germany and sponsored by an unrestricted educational grant from Novo Nordisk A/S, Bagsvård, Denmark. In the past 5 years, D.R.S. has received honoraria or travel support for consulting or lecturing from the following companies: Abbott AG, Baar, Switzerland, AstraZeneca AG, Zug, Switzerland, Bayer (Schweiz) AG, Zürich, Switzerland, Baxter S.p.A., Roma, Italy, B. Braun Melsungen AG, Melsungen, Germany, Boehringer Ingelheim (Schweiz) GmbH, Basel, Switzerland, Bristol-Myers-Squibb, Rueil-Malmaison Cedex, France, CSL Behring GmbH, Hattersheim am Main, Germany and Bern, Switzerland, Curacyte AG, Munich, Germany, Ethicon Biosurgery, Sommerville, New Jersey, USA, Fresenius SE, Bad Homburg v.d.H., Germany, Galenica AG, Bern, Switzerland (including Vifor SA, Villars-sur-Glâne, Switzerland), GlaxoSmithKline GmbH & Co. KG, Hamburg, Germany, Janssen-Cilag AG, Baar, Switzerland, Novo Nordisk A/S, Bagsvård, Denmark, Octapharma AG, Lachen, Switzerland, Organon AG, Pfäffikon/SZ, Switzerland, Oxygen Biotherapeutics, Costa Mesa, CA, Pentapharm GmbH (now tem Innovations GmbH), Munich, Germany, Roche Pharma (Schweiz) AG, Reinach, Switzerland and Schering-Plough International, Inc., Kenilworth, New Jersey, USA. P.S. has no conflicts of interest to declare.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Champion HR, Bellamy RF, Roberts CP, Leppaniemi A. A profile of combat injury. *J Trauma* 2003; 54:S13–S19.
2. Cothren CC, Moore EE, Hedegaard HB, Meng K. Epidemiology of urban trauma deaths: a comprehensive reassessment 10 years later. *World J Surg* 2007; 31:1507–1511.
3. Spahn DR, Bouillon B, Cerny V, *et al.* Management of bleeding and coagulopathy following major trauma: an updated European guideline. *Crit Care* 2013; 17:R76.

This article is the updated version of the European Trauma guideline published in 2007 and updated in 2010. Key changes include new recommendations on the appropriate use of vasopressors and inotropic agents, and reflect an awareness of the growing number of patients in the population at large treated with antiplatelet agents and/or oral anticoagulants. A significant addition is a new section that discusses the need for every institution to develop, implement and adhere to an evidence-based clinical protocol to manage traumatically injured patients. The remaining recommendations have been re-evaluated and graded based on literature published since the last edition of the guideline. Consideration was also given to the changes in clinical practice that have taken place during this time period as a result of both new evidence and changes in the general availability of relevant agents and technologies.

4. Esposito TJ, Sanddal TL, Reynolds SA, Sanddal ND. Effect of a voluntary trauma system on preventable death and inappropriate care in a rural state. *J Trauma* 2003; 54:663–670.

5. Rossaint R, Bouillon B, Cerny V, *et al.* The STOP the Bleeding Campaign. *Crit Care* 2013; 17:136.

This article deals with the fact that traumatic injuries worldwide are responsible for over 5 million deaths annually. Bleeding caused by traumatic injury-associated coagulopathy is the leading cause of potentially preventable death among trauma patients. Despite these facts, awareness of this problem is insufficient and treatment options are often unclear. The STOP the Bleeding Campaign therefore aims to increase awareness of this fact and its appropriate management by publishing European guidelines for the management of the bleeding trauma patient, by promoting and monitoring the implementation of these guidelines and by preparing promotional and educational material, organizing activities and developing health quality management tools. The campaign aims to reduce the number of patients who die within 24 h after arrival in the hospital due to exsanguination by a minimum of 20% within the next 5 years.

6. Kauvar DS, Lefering R, Wade CE. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma* 2006; 60:S3–11.
7. Maegele M, Lefering R, Yucel N, *et al.* Early coagulopathy in multiple injury: an analysis from the German Trauma Registry on 8724 patients. *Injury* 2007; 38:298–304.
8. MacLeod JB, Lynn M, McKenney MG, *et al.* Early coagulopathy predicts mortality in trauma. *J Trauma* 2003; 55:39–44.
9. Frith D, Goslings JC, Gaarder C, *et al.* Definition and drivers of acute traumatic coagulopathy: clinical and experimental investigations. *J Thromb Haemost* 2010; 8:1919–1925.
10. Brohi K, Singh J, Heron M, Coats T. Acute traumatic coagulopathy. *J Trauma* 2003; 54:1127–1130.
11. Brohi K, Cohen MJ, Ganter MT, *et al.* Acute traumatic coagulopathy: initiated by hypoperfusion: modulated through the protein C pathway? *Ann Surg* 2007; 245:812–818.
12. Carson JL, Carless PA, Hebert PC. Transfusion thresholds and other strategies for guiding allogeneic red blood cell transfusion. *Cochrane Database Syst Rev* 2012; 4:CD002042.

This Cochrane review shows that the existing evidence supports the use of restrictive transfusion triggers in most patients including those with preexisting cardiovascular disease. As there are no trials, the effects of restrictive transfusion triggers in high risk groups such as acute coronary syndrome need to be tested in further large clinical trials. In countries with inadequate screening of donor blood, the data may constitute a stronger basis for avoiding transfusion with allogeneic red cells.

13. Goodnough LT, Shander A. Patient blood management. *Anesthesiology* 2012; 116:1367–1376.

This article highlights the need for Patient Blood Management Programm and points out the fact that blood transfusions are potentially harmful and could also be used as quality indicators for healthcare institutions.

14. Shander A, Hofmann A, Isbister J, Van Aken H. Patient blood management: ■ the new frontier. *Best Pract Res Clin Anaesthesiol* 2013; 27:5–10.
- Blood transfusions face many issues including questionable safety and efficacy, increasing costs and limited supply. The need to provide effective care for a relatively small population of patients who could not be transfused for various reasons gave rise to 'bloodless medicine and surgery', which was subsequently proposed as a care strategy for all patients, with the goal of minimizing the use of allogeneic blood components. The next evolution came from the shift from a 'product-centred' approach towards a 'patient-centred' approach, that is, a focus on patient outcome rather than use of blood components, which gave birth to 'Patient Blood Management'.

15. Spahn DR, Goodnough LT. Alternatives to blood transfusion. *Lancet* 2013; ■ 381:1855–1865.

This article summarizes the current roles of alternatives to blood in the management of medical and surgical anaemias.

16. Williamson LM, Devine DV. Challenges in the management of the blood ■ supply. *Lancet* 2013; 381:1866–1875.

In the next 5–10 years, blood availability in developed countries will need to increase again to meet the demands of ageing populations. Increasing of the blood supply raises many challenges; new approaches to recruitment and retention of future generations of blood donors will be needed, and care will be necessary to avoid taking too much blood from these donors. Personalized medicine could be applied to match donors to patients, not only with extended blood typing, but also by using genetically determined storage characteristics of blood components. Growing of red cells or platelets in large quantities from stem cells is a possibility in the future, but challenges of cost, scaling up and reproducibility remain to be solved.

17. Park MS, Martini WZ, Dubick MA, *et al.* Thromboelastography as a better indicator of hypercoagulable state after injury than prothrombin time or activated partial thromboplastin time. *J Trauma* 2009; 67:266–276.
18. Davenport R, Manson J, De'Ath H, *et al.* Functional definition and characterization of acute traumatic coagulopathy. *Crit Care Med* 2011; 39:2652–2658.
19. Johansson PI, Stissing T, Bochen L, Ostrowski SR. Thromboelastography and tromboelastometry in assessing coagulopathy in trauma. *Scand J Trauma Resusc Emerg Med* 2009; 17:45.

20. Johansson PI, Stensballe J, Ostrowski SR. Current management of massive hemorrhage in trauma. *Scand J Trauma Resusc Emerg Med* 2012; 20:47.
 21. Johansson PI. Goal-directed hemostatic resuscitation for massively bleeding patients: the Copenhagen concept. *Transfus Apher Sci* 2010; 43:401–405.
 22. Ives C, Inaba K, Branco BC, *et al.* Hyperfibrinolysis elicited via thromboelastography predicts mortality in trauma. *J Am Coll Surg* 2012; 215:496–502.
- In this prospective analysis, hyperfibrinolysis on thromboelastography developed in approximately 10% of patients and was considerably more likely to require massive transfusion. Hyperfibrinolysis was a strong independent predictor of mortality. Additional evaluation of the role of thromboelastography-directed antifibrinolytic therapies is warranted.
23. Okamoto S, Hijikata-Okunomiya A, Wanaka K, *et al.* Enzyme-controlling medicines: introduction. *Semin Thromb Hemost* 1997; 23:493–501.
 24. Henry DA, Carless PA, Moxey AJ, *et al.* Antifibrinolytic use for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2011. doi: 10.1002/14651858. [Epub ahead of print]
 25. Shakur H, Roberts I, Bautista R, *et al.* Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet* 2010; 376:23–32.
 26. Grassin-Delyle S, Couturier R, Abe E, *et al.* A practical tranexamic acid dosing scheme based on population pharmacokinetics in children undergoing cardiac surgery. *Anesthesiology* 2013; 118:853–862.
- The authors report for the first time the pharmacokinetics of tranexamic acid in children undergoing cardiac surgery with CPB, and propose a dosing scheme for optimized tranexamic acid administration in those children.
27. Waldo T, Szlapka M, Haferkorn M, *et al.* Prospective clinical trial on dosage optimizing of tranexamic acid in nonemergency cardiac surgery procedures. *Clin Hemorheol Microcirc* 2013. doi: 10.3233/CH-131782. [Epub ahead of print]
 28. Ker K, Kiriya J, Perel P, *et al.* Avoidable mortality from giving tranexamic acid to bleeding trauma patients: an estimation based on WHO mortality data, a systematic literature review and data from the CRASH-2 trial. *BMC Emerg Med* 2012; 12:3.
- This study shows that the use of TXA in the treatment of traumatic bleeding has the potential to prevent many premature deaths every year. A large proportion of the potential health gains are in low and middle income countries.
29. Lipsky AM, Abramovich A, Nadler R, *et al.* Tranexamic acid in the prehospital setting: Israel Defense Forces' initial experience. *Injury* 2013; 45:66–70.
- The authors show that TXA may be successfully given in the prehospital setting without any apparent delays in evacuation. In light of recent evidence, the ability to give TXA closer to the time of wounding represents an important step toward improving the survival of trauma victims with haemorrhage, even before definitive care is available. TXA could be considered a viable option for use by advanced life support providers at or near the point of injury.
30. Roberts I, Shakur H, Afolabi A, *et al.* The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. *Lancet* 2011; 377:1096–1102.
 31. Roberts I, Shakur H, Coats T, *et al.* The CRASH-2 trial: a randomised controlled trial and economic evaluation of the effects of tranexamic acid on death, vascular occlusive events and transfusion requirement in bleeding trauma patients. *Health Technol Assess* 2013; 17:1–79.
- This analysis shows that the early administration of TXA safely reduced the risk of death in bleeding trauma patients and is highly cost effective. Treatment beyond 3 h of injury is unlikely to be effective.
32. Perel P, Ker K, Morales Uribe CH, Roberts I. Tranexamic acid for reducing mortality in emergency and urgent surgery. *Cochrane Database Syst Rev* 2013; 1:CD010245.
- The authors show by a prognostic model that one can obtain valid predictions of mortality in patients with traumatic bleeding. TXA can be administered safely to a wide spectrum of bleeding trauma patients and should not be restricted to the most severely injured. It has to be evaluated whether or not this model used in clinical practice has an impact on the management and outcomes of trauma patients.
33. Bhananker SM, Ramaiah R. Trends in trauma transfusion. *Int J Crit Illn Inj Sci* 2011; 1:51–56.
 34. Nakstad AR, Skaga NO, Pillgram-Larsen J, *et al.* Trends in transfusion of trauma victims: evaluation of changes in clinical practice. *Scand J Trauma Resusc Emerg Med* 2011; 19:23.
 35. Theusinger OM, Madjdpour C, Spahn DR. Resuscitation and transfusion management in trauma patients: emerging concepts. *Curr Opin Crit Care* 2012; 18:661–670.
 36. Luddington RJ. Thrombelastography/thromboelastometry. *Clin Lab Haematol* 2005; 27:81–90.
 37. Ganter MT, Hofer CK. Coagulation monitoring: current techniques and clinical use of viscoelastic point-of-care coagulation devices. *Anesth Analg* 2008; 106:1366–1375.
 38. Clauss A. Rapid physiological coagulation method in determination of fibrinogen. *Acta Haematol* 1957; 17:237–246.
 39. de Araujo Azi LM, Lopes FM, Garcia LV. Postoperative management of severe acute anemia in a Jehovah's Witness. *Transfusion* 2013; 11:142–143.
 40. Lauscher P, Kertscho H, Schmidt O, *et al.* Determination of organ-specific anemia tolerance. *Crit Care Med* 2013; 41:1037–1045.
- In this trial, pigs were randomized into three different groups and hemodiluted with hydroxyethyl starch comparing no hemodilution vs. hemoglobin of 4.0 g/dl and vs. the critical hemoglobin level of 2.7 g/dl. In the hemodiluted state 10 mg/kg of pimonidazole was injected, which forms protein adducts in hypoxic cells. Interestingly, metabolic parameters and oxygen consumption did not show that tissue oxygenation was restricted before reaching a hemoglobin level of 2.7 g/dl. Kidneys and skeletal muscle showed enhanced pimonidazole binding and vascular endothelial growth factor expression at a hemoglobin level of 4 g/dl. Other organs like heart, brain and liver showed no signs of tissue hypoxia at hemoglobin levels of 4 g/dl.
41. Pape A, Kertscho H, Stein P, *et al.* Neuromuscular blockade with rocuronium bromide increases the tolerance of acute normovolemic anemia in anesthetized pigs. *Eur Surg Res* 2012; 48:16–25.
- Deep neuromuscular blockade increases the tolerance of acute normovolemic anemia. The underlying mechanism most likely involves a reduction of skeletal muscular oxygen consumption. During acellular treatment of an acute blood loss, neuromuscular blockade might play an adjuvant role in situations in which profound stages of normovolemic anemia have to be tolerated.
42. Spahn DR, Rossaint R. Coagulopathy and blood component transfusion in trauma. *Br J Anaesth* 2005; 95:130–139.
 43. Mahambrey T, Pendry K, Nee A, *et al.* Critical care in emergency department: massive haemorrhage in trauma. *Emerg Med J* 2013; 30:9–14.
- This article addresses current concepts in haemostatic resuscitation. Recent guidelines on the diagnosis and treatment of coagulopathy in major trauma, and the role of component and adjuvant therapies, are considered. Finally, the potential role of thromboelastography and ROTEM are discussed.
44. Kozek-Langenecker S, Sorensen B, Hess JR, Spahn DR. Clinical effectiveness of fresh frozen plasma compared with fibrinogen concentrate: a systematic review. *Crit Care* 2011; 15:R239.
 45. Rossaint R, Bouillon B, Cerny V, *et al.* Management of bleeding following major trauma: an updated European guideline. *Crit Care* 2010; 14:R52.
 46. Schochl H, Nienaber U, Maegele M, *et al.* Transfusion in trauma: thromboelastometry-guided coagulation factor concentrate-based therapy versus standard fresh frozen plasma-based therapy. *Crit Care* 2011; 15:R83.
 47. Schochl H, Nienaber U, Hofer G, *et al.* Goal-directed coagulation management of major trauma patients using thromboelastometry (ROTEM)-guided administration of fibrinogen concentrate and prothrombin complex concentrate. *Crit Care* 2010; 14:R55.
 48. Theusinger OM, Felix C, Spahn DR. Strategies to reduce the use of blood products: a European perspective. *Curr Opin Anaesthesiol* 2012; 25:59–65.
 49. Theusinger OM, Baulig W, Seifert B, *et al.* Relative concentrations of haemostatic factors and cytokines in solvent/detergent-treated and fresh-frozen plasma. *Br J Anaesth* 2011; 106:505–511.
 50. Theusinger OM, Spahn DR, Ganter MT. Transfusion in trauma: why and how should we change our current practice? *Curr Opin Anaesthesiol* 2009; 22:305–312.
 51. Godej O, Gallmeier U, Schelian M, *et al.* Coagulation factor XIII reduces postoperative bleeding after coronary surgery with extracorporeal circulation. *Thorac Cardiovasc Surg* 2006; 54:26–33.
 52. Theusinger OM. The inhibiting effect of factor XIII on hyperfibrinolysis. *Anesth Analg* 2012; 114:1149–1150.
 53. Gorlinger K, Dirkmann D, Hanke AA, *et al.* First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: a retrospective, single-center cohort study. *Anesthesiology* 2011; 115:1179–1191.
 54. Sorensen B, Spahn DR, Innerhofer P, *et al.* Clinical review: prothrombin complex concentrates – evaluation of safety and thrombogenicity. *Crit Care* 2011; 15:201.
 55. Preston FE, Laidlaw ST, Sampson B, Kitchen S. Rapid reversal of oral anticoagulation with warfarin by a prothrombin complex concentrate (Beriplex): efficacy and safety in 42 patients. *Br J Haematol* 2002; 116:619–624.
 56. Nienaber U, Innerhofer P, Westermann I, *et al.* The impact of fresh frozen plasma vs coagulation factor concentrates on morbidity and mortality in trauma-associated haemorrhage and massive transfusion. *Injury* 2011; 42:697–701.
 57. Pape H-C, Neugebauer E, Ridley SA, *et al.* Cost-drivers in acute treatment of severe trauma in Europe: a systematic review of literature. *Eur J Trauma Emerg Surg* 2009; 35:61–66.
 58. Ruggeri ZM, Mannucci PM, Lombardi R, *et al.* Multimeric composition of factor VIII/von Willebrand factor following administration of DDAVP: implications for pathophysiology and therapy of von Willebrand's disease subtypes. *Blood* 1982; 59:1272–1278.
 59. Castaman G, Goodeve A, Eikenboom J. Principles of care for the diagnosis and treatment of von Willebrand disease. *Haematologica* 2013; 98:667–674.
 60. Beynon C, Sakowitz OW, Unterberg AW. Multiple electrode aggregometry in antiplatelet-related intracerebral haemorrhage. *J Clin Neurosci* 2013; 20:1805–1806.

61. Hansson EC, Shams Hakimi C, Astrom-Olsson K, *et al.* Effects of ex vivo platelet supplementation on platelet aggregability in blood samples from patients treated with acetylsalicylic acid, clopidogrel, or ticagrelor. *Br J Anaesth* 2013. doi: 10.1093/bja/aet333. [Epub ahead of print]

The authors investigated platelet aggregability with multiple electrode aggregometry with adenosine diphosphate, arachidonic acid and thrombin receptor activating peptide-6 as activators in whole-blood samples from patients treated with ASA, ASA + clopidogrel or ASA + ticagrelor, and from healthy controls. Aggregability was measured before and after supplementation of ABO-compatible fresh apheresis platelets. Platelet supplementation improved platelet aggregability independently of antiplatelet therapy. The effect on ADP-dependent platelet inhibition was limited however. Reduced effect of platelet transfusion is more likely within 2 h of drug intake in patients treated with ASA + ticagrelor compared with ASA + clopidogrel.

62. Croft PE, Cabral KP, Strout TD, *et al.* Managing blunt trauma in patients receiving dabigatran etexilate: case study and review of the literature. *J Emerg Nurs* 2013; 39:302–308.

This case presented by the authors shows that the rapid aging of the population increases the likelihood that injured elderly patients will have anticoagulation. The advantages of using dabigatran etexilate instead of warfarin will likely drive widespread use. These phenomena will certainly increase the probability that emergency clinicians will manage an injured patient taking the drug. Acute reversal of the effects of dabigatran etexilate in the actively bleeding patient is a significant challenge. Conventional coagulation studies provide an imperfect assessment of physiology, and data on effective reversal protocols are limited.

63. Wassef SN, Abel TJ, Grossbach A, *et al.* Traumatic intracranial hemorrhage in patients taking dabigatran: report of 3 cases and review of the literature. *Neurosurgery* 2013; 73:E368–E374.

The authors present their experience with three patients who suffered an intracranial hemorrhage either spontaneously or after low-energy cranial trauma and review the available literature describing dabigatran use in patients with traumatic brain injury. Intracranial hemorrhage in patients taking anticoagulants and/or antiplatelets can have either a benign or malignant clinical course. At this time, there is little experience with dabigatran reversal; however, several strategies for rapid reversal have been proposed. All patients with intracranial hemorrhage taking dabigatran should be admitted for close neurological monitoring and serial imaging.

64. Simonsen CZ, Steiner T, Tietze A, Damgaard D. Dabigatran-related intracerebral hemorrhage resulting in hematoma expansion. *J Stroke Cerebrovasc Dis* 2013. doi: 10.1016/j.jstrokecerebrovasdis.2013.08.011. [Epub ahead of print]

The authors present a case that suggests that hematoma expansion may occur after novel oral anticoagulants-related intracerebral hemorrhage.

65. Schulman S, Kearon C, Kakkar AK, *et al.* Extended use of dabigatran, warfarin, or placebo in venous thromboembolism. *N Engl J Med* 2013; 368:709–718.

This study showed that dabigatran is effective in the extended treatment of venous thromboembolism and carried a lower risk of major or clinically relevant bleeding than warfarin but a higher risk than placebo.

66. Eerenberg ES, Kamphuisen PW, Sijpkens MK, *et al.* Reversal of rivaroxaban and dabigatran by prothrombin complex concentrate: a randomized, placebo-controlled, crossover study in healthy subjects. *Circulation* 2011; 124:1573–1579.

67. Davis PK, Musunuru H, Walsh M, *et al.* The ex vivo reversibility of dabigatran-induced whole-blood coagulopathy as monitored by thromboelastography: mechanistic implications for clinical medicine. *Thromb Haemost* 2012; 108:586–588.

68. Marlu R, Hodaj E, Paris A, *et al.* Effect of nonspecific reversal agents on anticoagulant activity of dabigatran and rivaroxaban: a randomised crossover ex vivo study in healthy volunteers. *Thromb Haemost* 2012; 108:217–224.

The new anticoagulants dabigatran and rivaroxaban are responsible for haemorrhagic complications. The authors tested the effect of all putative haemostatic agents on the anticoagulant activity of these new drugs using thrombin generation tests. In an ex-vivo study, 10 healthy white male individuals were randomized to receive rivaroxaban (20 mg) or dabigatran (150 mg) in one oral administration. After a 2 weeks washout period, they received the other anticoagulant. Venous blood samples were collected just before drug administration and 2 h thereafter. Reversal of anticoagulation was tested *in vitro* using PCC, rFVIIa or FEIBA(R) at various concentrations. PCC strongly corrected endogenous thrombin potential-area under the curve (ETP-AUC), whereas rFVIIa only modified the kinetic parameters. FEIBA corrected all parameters. Dabigatran specially affects the kinetics of thrombin generation with prolonged lag-time and time to peak. Although PCC increased ETP-AUC, only rFVIIa and FEIBA corrected the altered lag time. For both anticoagulants, lower doses of FEIBA, corresponding to a quarter to half the dose usually used, have potential reversal profile of interest. Some nonspecific reversal agents appear to be able to reverse the anticoagulant activity of rivaroxaban or dabigatran.

69. Warkentin TE, Margetts P, Connolly SJ, *et al.* Recombinant factor VIIa (rFVIIa) and hemodialysis to manage massive dabigatran-associated postcardiac surgery bleeding. *Blood* 2012; 119:2172–2174.

70. Singh T, Maw TT, Henry BL, *et al.* Extracorporeal therapy for dabigatran removal in the treatment of acute bleeding: a single center experience. *Clin J Am Soc Nephrol* 2013; 8:1533–1539.

This study shows that dabigatran concentrations decreased during intermittent hemodialysis but rebounded up within 2 h after the completion of dialysis. Initiation of continuous renal replacement therapy after intermittent hemodialysis attenuated the rebound effect in one patient and contributed to a reduction in dabigatran concentrations of 81% over 30 h. Extracorporeal therapy lowered dabigatran concentrations, suggesting that it removed the drug and may effectively accelerate total clearance, specially in patients with impaired kidney function. The use of prolonged intermittent hemodialysis or intermittent hemodialysis followed by continuous renal replacement therapy is recommended for the management of life-threatening bleeding in patients receiving dabigatran.

71. Wanek MR, Horn ET, Elapavaluru S, *et al.* Safe use of hemodialysis for dabigatran removal before cardiac surgery. *Ann Pharmacother* 2012; 46:e21.

72. Chang DN, Dager WE, Chin AI. Removal of dabigatran by hemodialysis. *Am J Kidney Dis* 2013; 61:487–489.

73. van Ryn J, Stangier J, Haertter S, *et al.* Dabigatran etexilate: a novel, reversible, oral direct thrombin inhibitor – interpretation of coagulation assays and reversal of anticoagulant activity. *Thromb Haemost* 2010; 103:1116–1127.

74. Lu G, DeGuzman FR, Hollenbach SJ, *et al.* A specific antidote for reversal of anticoagulation by direct and indirect inhibitors of coagulation factor Xa. *Nat Med* 2013; 19:446–451.